

The Complexity of Subsumption in Fuzzy \mathcal{EL}

Stefan Borgwardt

Chair for Automata Theory
Technische Universität Dresden
Germany

Stefan.Borgwardt@tu-dresden.de

Marco Cerami

Department of Computer Science
Palacký University in Olomouc
Czech Republic

marco.cerami@upol.cz

Rafael Peñaloza

KRDB Research Centre
Free University of Bozen-Bolzano
Italy

rafael.penaloz@unibz.it

Abstract

Fuzzy Description Logics (DLs) are used to represent and reason about vague and imprecise knowledge that is inherent to many application domains. It was recently shown that the complexity of reasoning in finitely valued fuzzy DLs is often not higher than that of the underlying classical DL. We show that this does not hold for fuzzy extensions of the light-weight DL \mathcal{EL} , which is used in many biomedical ontologies, under the Łukasiewicz semantics. The complexity of reasoning increases from PTIME to EXPTIME, even if only one additional truth value is introduced. The same lower bound holds also for infinitely valued Łukasiewicz extensions of \mathcal{EL} .

1 Introduction

Description Logics (DLs) are a family of knowledge representation formalisms that are successfully applied in many application domains. They provide the logical foundation for the Direct Semantics of the standard web ontology language OWL 2.¹ The light-weight DL \mathcal{EL} , underlying the OWL 2 EL profile, is of particular interest since all common reasoning problems are polynomial in this logic, and it is used in many prominent biomedical ontologies like SNOMED CT² and the Gene Ontology.³ Knowledge is represented by a set of *general concept inclusions (GCIs)* like

$$\exists \text{hasDisease.Flu} \sqsubseteq \exists \text{hasSymptom.Headache} \sqcap \exists \text{hasSymptom.Fever} \quad (1)$$

which states that every patient with a flu must also show headache and fever as symptoms. Reasoning in \mathcal{EL} is a polynomial problem [Baader *et al.*, 2005].

An important problem for AI practical applications is to represent and reason with vague or imprecise knowledge in a formal way. Fuzzy Description Logics (FDLs) [Straccia, 2001; Hájek, 2005] were introduced with this goal in mind. The main premise of fuzzy logics is the use of more than

two truth degrees to allow a more fine-grained analysis of dependencies between concepts. Usually, these degrees are arranged in a totally ordered algebra, or *chain*, in the interval $[0, 1]$. A patient having a body temperature of 37.5°C can have a degree of fever of 0.5, whereas a temperature of 39.2°C may be interpreted as a fever with degree of 0.9. Considering the GCI (1), the severity of the symptoms certainly influences the severity of the disease, and thus truth degrees can be transferred between concepts. Depending on the granularity one wants to have, one can choose to allow 10 or 100 truth degrees, or even admit the whole interval $[0, 1]$. Another degree of freedom in FDLs comes from the choice of possible semantics for the logical constructors. The most general semantics are based on *triangular norms (t-norms)* that are used to interpret conjunctions. Among these, the most prominent ones are the *Gödel*, *Łukasiewicz*, and *product* t-norms. All (continuous) t-norms over chains can be expressed as combinations of these three basic ones.

Unfortunately, reasoning in many infinitely valued FDLs becomes undecidable [Baader and Peñaloza, 2011; Cerami and Straccia, 2013]. For a systematic study on this topic, see [Borgwardt *et al.*, 2015b]. On the other hand, every finitely valued FDL that has been recently studied has not only been proved to be decidable, but even to belong to the same complexity class as the corresponding classical DL [Borgwardt and Peñaloza, 2013; 2014; Bou *et al.*, 2012].

A question that naturally arises is whether the finitely valued fuzzy framework always yields the same computational complexity as the corresponding classical formalisms. A common opinion is that everything that can be expressed in finitely valued FDLs can be reduced to the corresponding classical DLs without any serious loss of efficiency. Indeed, although some known direct translations of finitely valued FDLs into classical DLs are exponential [Bobillo and Straccia, 2011], more efficient reasoning can be achieved through direct algorithms [Borgwardt and Peñaloza, 2013]. The problem of finding a complexity gap between classical and finitely valued logics has already been considered. In [Cerami and Straccia, 2014], the authors analyze different constructors that could cause an increase in the complexity, but no specific answer is found. In [Borgwardt *et al.*, 2014] it is shown that the Łukasiewicz t-norm is a source of nondeterminism able to cause a significant increase in expressivity in very simple propositional languages. In this work, we build on the meth-

¹<http://www.w3.org/TR/owl2-overview/>

²<http://www.ihtsdo.org/snomed-ct/>

³<http://geneontology.org/>

ods of [Borgwardt *et al.*, 2014] to show even more dramatic increases in complexity for finitely valued extensions of \mathcal{EL} .

The question about the computational complexity of \mathcal{EL} under infinitely valued semantics has been already considered. Borgwardt and Peñaloza show that reasoning in \mathcal{EL} under semantics including the Łukasiewicz t-norm is CO-NP-hard [2013], but the proof does not apply to the finitely valued case. In contrast, infinitely valued Gödel semantics do not increase the complexity of reasoning [Mailis *et al.*, 2012].

In this work, we prove that \mathcal{EL} under finitely valued semantics is EXPTIME-complete whenever the Łukasiewicz t-norm is included in the semantics. This proves a dichotomy similar to one that exists for infinitely valued FDLs [Borgwardt *et al.*, 2015b] since, for all other finitely valued chains of truth values, reasoning in fuzzy \mathcal{EL} can be shown to be in PTIME using the methods from [Mailis *et al.*, 2012]. The relevance of our result goes beyond the computational aspect. Indeed, this is so far the first instance of a finitely valued DL that is more complex than the same language under classical semantics. In this way, we obtain an answer to the open problem whether finitely valued FDLs and classical DLs are equally powerful, at least from a computational complexity point of view. As a side benefit, we obtain the same (EXPTIME) lower bound for the complexity of infinitely valued fuzzy extensions of \mathcal{EL} that use the Łukasiewicz t-norm, improving the lower bound from [Borgwardt and Peñaloza, 2013]. Full proofs of all results appear in the technical report [Borgwardt *et al.*, 2015a].

2 Preliminaries

Fuzzy DLs extend classical DLs by allowing more than two truth degrees in the semantics. We first introduce the classes of truth degrees relevant for this paper and then recall the logics \mathcal{ELU} and $\mathcal{L}\text{-}\mathcal{EL}$.

2.1 Chains of Truth Values

A *chain* is an algebra \mathbf{L} whose elements are linearly ordered; that is, for every $x, y \in \mathbf{L}$ it holds that either $x \leq y$ or $y \leq x$. We consider here chains of the form $(\mathbf{L}, *_\mathbf{L}, \Rightarrow_\mathbf{L})$, where

- \mathbf{L} is a subset of $[0, 1]$ that contains 0 and 1;
- the *t-norm* $*_\mathbf{L}$ is a binary operator on \mathbf{L} that is associative, commutative, monotone, and has 1 as unit element; and
- the *residuum* $\Rightarrow_\mathbf{L}$ of $*_\mathbf{L}$ is a binary operator on \mathbf{L} that satisfies $(x *_\mathbf{L} y) \leq z$ iff $y \leq (x \Rightarrow_\mathbf{L} z)$ for all $x, y, z \in \mathbf{L}$.

An element $x \in \mathbf{L}$ is *idempotent* if $x *_\mathbf{L} x = x$. For ease of presentation, we will often identify \mathbf{L} and $(\mathbf{L}, *_\mathbf{L}, \Rightarrow_\mathbf{L})$ and omit the subscript \mathbf{L} if the chain we use is clear from the context.

We consider in particular the two cases where (i) \mathbf{L} is defined over the interval $[0, 1]$ of real numbers, or (ii) \mathbf{L} is a *finite chain*. In the former case, we always make the assumption that the operator $*_\mathbf{L}$ is *continuous* as a function from $[0, 1] \times [0, 1]$ to $[0, 1]$. One reason for this assumption is that it ensures that the residuum is uniquely determined by the t-norm [Klement *et al.*, 2000]. In case (ii), we similarly assume that $*_\mathbf{L}$ is *smooth*, i.e. for every $x, y, z \in \mathbf{L}$, whenever x and y are direct neighbors in \mathbf{L} , with $x < y$, then there is no $w \in \mathbf{L}$ such that $x *_\mathbf{L} z < w < y *_\mathbf{L} z$ [Mayor and

Table 1: Basic t-norms and their residua

	t-norm $x * y$	residuum $x \Rightarrow y$
$\mathbf{G}_n / \mathbf{G}$	$\min(x, y)$	$\begin{cases} 1 & \text{if } x \leq y \\ y & \text{if } x > y \end{cases}$
$\mathbf{L}_n / \mathbf{L}$	$\max(x + y - 1, 0)$	$\min(1 - x + y, 1)$
Π	$x \cdot y$	$\begin{cases} 1 & \text{if } x \leq y \\ \frac{y}{x} & \text{if } x > y \end{cases}$

Torrens, 1993]. If $*_\mathbf{L}$ is continuous (smooth), then we call \mathbf{L} *continuous (smooth)*.

By restricting the algebra of truth values to two elements, the classical Boolean algebra of truth and falsity $\mathbf{B} = (\{0, 1\}, *_\mathbf{B}, \Rightarrow_\mathbf{B})$ is obtained. In this case, $*_\mathbf{B}$ and $\Rightarrow_\mathbf{B}$ are the classical conjunction and the material implication, respectively.

The most interesting chains with continuous or smooth t-norms are the ones given by the Gödel (\mathbf{G}), Łukasiewicz (\mathbf{L}), and product (Π) t-norms. The finitely valued versions of the former two, denoted by \mathbf{L}_n and \mathbf{G}_n for $n \geq 2$, are defined over the n -element chain $0 < \frac{1}{n-1} < \dots < \frac{n-2}{n-1} < 1$. These operators and their residua are defined in Table 1. Notice that a finitely valued version of the product t-norm Π cannot exist: the chain \mathbf{L} needs to be closed under the t-norm, but for any $x \in (0, 1)$, the set $\{x^m \mid m \geq 0\}$ is infinite.

The following properties, which are crucial for our reductions, follow directly from the previous definitions.

Fact 1. For all $x, y \in \mathbf{L}$ and $T \subseteq \mathbf{L}$, it holds that

- $x \Rightarrow_\mathbf{L} y = 1$ iff $x \leq y$;
- $1 \Rightarrow_\mathbf{L} x = x$;
- $x \Rightarrow_\mathbf{L} y \geq y$;
- if $\mathbf{L} = \mathbf{L}_n$, then $x *_\mathbf{L}_n y \geq \frac{n-2}{n-1}$ iff either $x = 1$ or $y = 1$;
- if $\mathbf{L} = \mathbf{L}_n$ and $x < 1$, then for all $m \geq n - 1$ we have $\underbrace{x *_\mathbf{L}_n \dots *_\mathbf{L}_n}_m x = 0$.

The t-norms defined so far can be used to build all other continuous t-norms over $[0, 1]$, and all smooth t-norms over finite chains, using the following construction.

Definition 2. Let \mathbf{L} be a chain, $(\mathbf{L}_i)_{i \in I}$ be a family of chains, and $(\lambda_i)_{i \in I}$ be isomorphisms between intervals $[a_i, b_i] \subseteq \mathbf{L}$ and \mathbf{L}_i such that the intersection of any two intervals contains at most one element. \mathbf{L} is the *ordinal sum* of the family $(\mathbf{L}_i, \lambda_i)_{i \in I}$ if, for all $x, y \in \mathbf{L}$,

$$x *_\mathbf{L} y = \begin{cases} \lambda_i^{-1}(\lambda_i(x) *_\mathbf{L}_i \lambda_i(y)) & \text{if } x, y \in (a_i, b_i), \\ \min\{x, y\} & \text{otherwise.} \end{cases}$$

Intuitively, the ordinal sum of the chains \mathbf{L}_i is a chain whose domain is built up by appending the domains of the chains \mathbf{L}_i and whose operation $*_\mathbf{L}$ is $*_{\mathbf{L}_i}$ when the operands belong to the same chain \mathbf{L}_i and it is \min otherwise.

Every chain over $[0, 1]$ with a continuous t-norm is isomorphic to an ordinal sum of infinitely valued Łukasiewicz

and product chains [Hájek, 2001; Mostert and Shields, 1957]. Similarly, every smooth finite chain is an ordinal sum of chains of the form \mathfrak{L}_n with $n \geq 3$ [Mayor and Torrens, 2005]. All elements that are not contained strictly within one such Łukasiewicz or product component are idempotent and can be thought of as belonging to a (finite or infinite) Gödel chain. We say that a (finite or infinite) chain *contains* the Łukasiewicz t-norm if its ordinal sum representation contains at least one Łukasiewicz component; similarly, it *starts with* the Łukasiewicz t-norm if it contains a Łukasiewicz component in an interval $[0, b]$. Note that every chain that contains the Łukasiewicz t-norm can be represented as the ordinal sum of an arbitrary chain \mathfrak{L}_1 and another chain \mathfrak{L}_2 that starts with the Łukasiewicz t-norm.

Another way to view these characterizations is to observe that every smooth finite chain is either a Gödel chain or contains at least one Łukasiewicz component, and every continuous chain over $[0, 1]$ is either a Gödel chain or contains at least one Łukasiewicz or product component. As we will see, analyzing the properties of the basic t-norms from Table 1 provides an insight into the general case, with arbitrary (continuous or smooth) t-norms.

2.2 \mathcal{ELU} and $\mathcal{L}\text{-}\mathcal{EL}$

A *description signature* is a tuple $(\mathbb{N}_C, \mathbb{N}_R)$, where \mathbb{N}_C and \mathbb{N}_R are disjoint countable sets of *concept names* and *role names*, respectively. \mathcal{EL} concepts are built inductively from concept and role names through the grammar rule $C, D ::= A \mid \top \mid C \sqcap D \mid \exists r.C$ where $A \in \mathbb{N}_C$ and $r \in \mathbb{N}_R$. \mathcal{ELU} concepts are formed by adding the option $C \sqcup D$ to the previous rule. In the rest of the paper we will use the abbreviation C^m , $m \geq 1$, for the m -ary conjunction; i.e. $C^1 := C$ and $C^{m+1} := C^m \sqcap C$.

There is often no difference between the syntax of classical and fuzzy languages. The differences between both frameworks begin when the *semantics* of concepts and roles is introduced. As remarked in Section 2.1, it suffices to restrict the semantics to the two-element chain \mathbf{B} to obtain the classical semantics. However, we define both semantics to aid understanding and readability of the proofs.

Fuzzy Semantics of $\mathcal{L}\text{-}\mathcal{EL}$.

Consider an arbitrary but fixed chain $\mathbf{L} = (\mathbf{L}, *, \Rightarrow)$. An \mathbf{L} -*interpretation* is a pair $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$ consisting of:

- a nonempty (classical) set $\Delta^{\mathcal{I}}$ (called *domain*), and
- a *fuzzy interpretation function* $\cdot^{\mathcal{I}}$ that assigns
 - to each $A \in \mathbb{N}_C$ a fuzzy set $A^{\mathcal{I}}: \Delta^{\mathcal{I}} \rightarrow \mathbf{L}$, and
 - to each $r \in \mathbb{N}_R$ a fuzzy relation $r^{\mathcal{I}}: \Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}} \rightarrow \mathbf{L}$.

The interpretation function is extended to \mathcal{EL} concepts inductively by defining, for all $x \in \mathbf{L}$,

$$\begin{aligned} \top^{\mathcal{I}}(x) &:= 1, \\ (C \sqcap D)^{\mathcal{I}}(x) &:= C^{\mathcal{I}}(x) * D^{\mathcal{I}}(x), \\ (\exists r.C)^{\mathcal{I}}(x) &:= \sup_{y \in \Delta^{\mathcal{I}}} r^{\mathcal{I}}(x, y) * C^{\mathcal{I}}(y). \end{aligned}$$

Classical semantics of \mathcal{ELU} .

A *classical interpretation* is a pair $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$, where

- $\Delta^{\mathcal{I}}$ is a nonempty (classical) set (called *domain*), and
- $\cdot^{\mathcal{I}}$ is an *interpretation function* that assigns:
 - to each $A \in \mathbb{N}_C$ a set $A^{\mathcal{I}} \subseteq \Delta^{\mathcal{I}}$, and
 - to each $r \in \mathbb{N}_R$ a binary relation $r^{\mathcal{I}} \subseteq \Delta^{\mathcal{I}} \times \Delta^{\mathcal{I}}$.

This function is extended to \mathcal{ELU} concepts by setting

$$\begin{aligned} \top^{\mathcal{I}} &:= \Delta^{\mathcal{I}}, \\ (C \sqcap D)^{\mathcal{I}} &:= C^{\mathcal{I}} \cap D^{\mathcal{I}}, \\ (C \sqcup D)^{\mathcal{I}} &:= C^{\mathcal{I}} \cup D^{\mathcal{I}}, \\ (\exists r.C)^{\mathcal{I}} &:= \{x \in \Delta^{\mathcal{I}} \mid \exists y \in \Delta^{\mathcal{I}}: (x, y) \in r^{\mathcal{I}} \wedge y \in C^{\mathcal{I}}\}. \end{aligned}$$

Clearly, by replacing the relation \in by its characteristic function $\chi_{\in}: \Delta^{\mathcal{I}} \rightarrow \{0, 1\}$, we obtain a special case of fuzzy semantics. Whenever \mathbf{L} is one of the specific chains introduced in the previous section, e.g. \mathfrak{L}_n , then we denote the resulting logic by $\mathfrak{L}_n\text{-}\mathcal{EL}$ instead of $\mathbf{L}\text{-}\mathcal{EL}$.

In infinite chains, interpretations are often restricted to be *witnessed* [Hájek, 2005], which means that for every existential restriction $\exists r.C$ and $x \in \Delta^{\mathcal{I}}$ there is an element $y \in \Delta^{\mathcal{I}}$ that realizes the supremum in the semantics of $\exists r.C$ at x , i.e. $(\exists r.C)^{\mathcal{I}}(x) = r^{\mathcal{I}}(x, y) * C^{\mathcal{I}}(y)$. Under finitely valued and classical semantics this property is always satisfied, and it corresponds to the intuition that an existential restriction actually forces the existence of a single domain element that satisfies it, instead of infinitely many that only satisfy the restriction in the limit. We also adopt the restriction to witnessed interpretations in what follows.

In DLs, the domain knowledge is represented by axioms that restrict the class of interpretations under consideration. In the fuzzy framework, these axioms are assigned a minimum degree of truth to which they must be satisfied. Graded *general concept inclusions* (GCIs) are expressions of the form $\langle C \sqsubseteq D \geq \ell \rangle$, where $\ell \in \mathbf{L}$. The \mathbf{L} -interpretation \mathcal{I} *satisfies* this axiom if $C^{\mathcal{I}}(x) \Rightarrow D^{\mathcal{I}}(x) \geq \ell$ holds for all $x \in \Delta^{\mathcal{I}}$. As usual, a TBox is a finite set of GCIs, and an \mathbf{L} -interpretation \mathcal{I} *satisfies* a TBox if it satisfies every axiom in it.

We consider the problem of deciding whether a concept C is ℓ -*subsumed* by another concept D with respect to a TBox \mathcal{T} for a value $\ell \in \mathbf{L} \setminus \{0\}$. That is, whether every \mathbf{L} -interpretation \mathcal{I} that satisfies \mathcal{T} also satisfies $\langle C \sqsubseteq D \geq \ell \rangle$. In the classical case, we talk simply about *subsumption*, and for $\ell = 1$ the problem simplifies to the question whether $C^{\mathcal{I}} \subseteq D^{\mathcal{I}}$ holds in all interpretations \mathcal{I} that satisfy \mathcal{T} .

In the particular case of $\mathbf{G}_n\text{-}\mathcal{EL}$, subsumption is decidable in polynomial time. This can be shown by generalizing the proof from [Baader *et al.*, 2005] (for $n = 2$), as it was done in [Mailis *et al.*, 2012] for the infinitely valued $\mathbf{G}\text{-}\mathcal{EL}$.

Proposition 3. *Deciding ℓ -subsumption with respect to a TBox in $\mathbf{G}_n\text{-}\mathcal{EL}$ is PTIME-complete.*

In this paper, we show that for all other finite chains the subsumption problem becomes EXPTIME-complete. As a first step, we show that this problem is EXPTIME-hard for all finite Łukasiewicz chains with at least three elements, and then use this result in Section 4 to show EXPTIME-hardness




Figure 1: Illustration of the reductions

under any finite chain that contains a Łukasiewicz component, i.e. is not of the form \mathbf{G}_n (see Definition 2). A matching EXPTIME upper bound was shown in [Borgwardt and Peñaloza, 2013]. In Section 5, we adapt our reduction to show EXPTIME-hardness of $\mathbf{L}\text{-}\mathcal{EL}$, and even for every continuous chain over $[0, 1]$ containing a Łukasiewicz component.

The idea behind the reductions is illustrated in Figure 1 for chains \mathbf{L} containing either an \mathbf{L}_3 -component or an infinitely valued \mathbf{L} -component. To simulate the semantics of \mathcal{ELU} , the values 0.5 and 1 in $\mathbf{L}_3\text{-}\mathcal{EL}$ (or $\mathbf{L}\text{-}\mathcal{EL}$) are used to simulate the truth values *false* and *true*, respectively. The chain \mathbf{L}_3 (\mathbf{L}) is then embedded into \mathbf{L} as depicted.

3 Finite Łukasiewicz Chains

We now reduce the subsumption problem of the classical DL \mathcal{ELU} to the subsumption problem of $\mathbf{L}_n\text{-}\mathcal{EL}$, where $n \geq 3$. Since concept subsumption in \mathcal{ELU} is EXPTIME-complete [Baader *et al.*, 2005], this reduction shows that the subsumption problem is EXPTIME-hard already for $\mathbf{L}_3\text{-}\mathcal{EL}$; i.e. for Łukasiewicz chains containing three truth degrees.

Note that it suffices to consider subsumption problems between two *concept names* since an \mathcal{ELU} concept C is subsumed by another \mathcal{ELU} concept D w.r.t. an \mathcal{ELU} TBox \mathcal{T} iff A is subsumed by B w.r.t. $\mathcal{T} \cup \{ \langle A \sqsubseteq C \rangle, \langle D \sqsubseteq B \rangle \}$, for two new concept names A, B [Baader *et al.*, 2005]. Furthermore, we can restrict our attention to \mathcal{ELU} TBoxes in *normal form*, which only contain axioms of the form

$$A_1 \sqcap A_2 \sqsubseteq B, \exists r.A \sqsubseteq B, A \sqsubseteq \exists r.B, \text{ or } A \sqsubseteq B_1 \sqcup B_2,$$

where A, A_1, A_2, B, B_1 and B_2 are concept names or \top . As shown in [Baader *et al.*, 2005], every \mathcal{ELU} TBox can be transformed in linear time into an equivalent one (w.r.t. the original signature) in normal form.

The main idea of our reduction is to simulate a classical concept name in $\mathbf{L}_n\text{-}\mathcal{EL}$ by considering all values below $\frac{n-2}{n-1}$ to be equivalent to 0, and thus only the value 1 can be used to express that a domain element belongs to the concept name. We can then express a classical disjunction of the form $B_1 \sqcup B_2$ by restricting the value of the fuzzy conjunction $B_1 \sqcap B_2$ to be $\geq \frac{n-2}{n-1}$: according to Fact 1, the latter is equivalent to B_1 or B_2 having value 1. Furthermore, we reformulate classical subsumption between C and D as

1-subsumption between C^{n-1} and D^{n-1} : again by Fact 1, the latter two concepts can take only the values 0 or 1. Notice that the conjunctions C^{n-1} and D^{n-1} are fundamental for this reduction to work; their purpose is to produce a crisp version of the concepts C and D .

More formally, let $n \geq 3$, \mathcal{T} be an \mathcal{ELU} TBox in normal form, and $C, D \in \mathbf{N}_C$. We construct an $\mathbf{L}_n\text{-}\mathcal{EL}$ TBox $\rho_n(\mathcal{T})$ such that C is subsumed by D w.r.t. \mathcal{T} if and only if C^{n-1} is 1-subsumed by D^{n-1} w.r.t. $\rho_n(\mathcal{T})$. Since \mathcal{T} is in normal form, we can define the reduction ρ_n for each of the four kinds of axioms listed above:

$$\begin{aligned} \rho_n(A_1 \sqcap A_2 \sqsubseteq B) &:= \langle A_1 \sqcap A_2 \sqsubseteq B \geq 1 \rangle \\ \rho_n(\exists r.A \sqsubseteq B) &:= \langle \exists r.A \sqsubseteq B \geq 1 \rangle \\ \rho_n(A \sqsubseteq \exists r.B) &:= \langle A \sqsubseteq (\exists r.B)^{n-1} \geq \frac{1}{n-1} \rangle \\ \rho_n(A \sqsubseteq B_1 \sqcup B_2) &:= \langle A \sqsubseteq B_1 \sqcap B_2 \geq \frac{n-2}{n-1} \rangle \end{aligned}$$

Finally, we set $\rho_n(\mathcal{T}) := \{ \rho_n(\alpha) \mid \alpha \in \mathcal{T} \}$. Notice that $\rho_n(\mathcal{T})$ has as many axioms as \mathcal{T} , and the size of each axiom is increased by a factor of at most n . Hence, the translation $\rho_n(\mathcal{T})$ can be performed in polynomial time. The translation of the axiom $A \sqsubseteq \exists r.B$ deserves special attention. Notice that $\rho_n(A \sqsubseteq \exists r.B)$ uses a conjunction of the concept $\exists r.B$ on the right-hand side. This is necessary to guarantee that we consider cases where both, the role relation, and the membership to B have degree 1. We show that this translation satisfies the properties described above.

First we show that if C is subsumed by D w.r.t. \mathcal{T} , then C^{n-1} is 1-subsumed by D^{n-1} w.r.t. $\rho_n(\mathcal{T})$. In order to achieve this result, for any \mathbf{L}_n -interpretation $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$ we define the classical interpretation $\mathcal{I}_{cr} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}_{cr}})$, where:

- $x \in A^{\mathcal{I}_{cr}}$ iff $A^{\mathcal{I}}(x) = 1$ for every $A \in \mathbf{N}_C$ and $x \in \Delta^{\mathcal{I}}$,
- $(x, y) \in r^{\mathcal{I}_{cr}}$ iff $r^{\mathcal{I}}(x, y) = 1$ for every $r \in \mathbf{N}_R$ and $x, y \in \Delta^{\mathcal{I}}$.

Recall that for every $x \in \Delta^{\mathcal{I}}$ it holds that $x \in \top^{\mathcal{I}_{cr}}$ and $\top^{\mathcal{I}}(x) = 1$. This means that \top behaves exactly like a concept name in this reduction.

It can be shown that if the \mathbf{L}_n -interpretation \mathcal{I} satisfies $\rho_n(E \sqsubseteq F)$ for some \mathcal{ELU} GCI $E \sqsubseteq F$ in normal form, then \mathcal{I}_{cr} satisfies $E \sqsubseteq F$.

Lemma 4. *Let \mathcal{I} be an \mathbf{L}_n -interpretation that satisfies $\rho_n(\mathcal{T})$. Then \mathcal{I}_{cr} satisfies \mathcal{T} .*

Proof Sketch. We consider only the two most interesting kinds of axioms here. Take first any $A \sqsubseteq \exists r.B \in \mathcal{T}$ and assume that $\langle A \sqsubseteq (\exists r.B)^{n-1} \geq \frac{1}{n-1} \rangle$ is satisfied by \mathcal{I} . For every element $x \in A^{\mathcal{I}_{cr}}$, we need to show that $x \in (\exists r.B)^{\mathcal{I}_{cr}}$. By the definition of \mathcal{I}_{cr} , we have $A^{\mathcal{I}}(x) = 1$. By our assumption, this implies that $((\exists r.B)^{n-1})^{\mathcal{I}}(x) \geq \frac{1}{n-1}$, and thus $(\exists r.B)^{\mathcal{I}}(x) = 1$ by Fact 1. Hence, $1 = \sup_{z \in \Delta^{\mathcal{I}}} r^{\mathcal{I}}(x, z) *_{\mathbf{L}_n} B^{\mathcal{I}}(z)$, i.e. there exists $y \in \Delta^{\mathcal{I}}$ such that $r^{\mathcal{I}}(x, y) = 1$ and $B^{\mathcal{I}}(y) = 1$. Again by the definition of \mathcal{I}_{cr} , we have $(x, y) \in r^{\mathcal{I}_{cr}}$ and $y \in B^{\mathcal{I}_{cr}}$, and hence $x \in (\exists r.B)^{\mathcal{I}_{cr}}$.

Consider now any $A \sqsubseteq B_1 \sqcup B_2 \in \mathcal{T}$ and assume that $\langle A \sqsubseteq B_1 \sqcap B_2 \geq \frac{n-2}{n-1} \rangle$ is satisfied by \mathcal{I} . If $x \in A^{\mathcal{I}_{cr}}$, then $A^{\mathcal{I}}(x) = 1$. By our assumption, this implies that

$(B_1 \sqcap B_2)^{\mathcal{I}}(x) \geq \frac{n-2}{n-1}$, and hence either $B_1^{\mathcal{I}}(x) = 1$ or $B_2^{\mathcal{I}}(x) = 1$ must hold by Fact 1. Again by the definition of \mathcal{I}_{cr} , we get that either $x \in B_1^{\mathcal{I}_{cr}}$ or $x \in B_2^{\mathcal{I}_{cr}}$. \square

Suppose now that C^{n-1} is not 1-subsumed by D^{n-1} w.r.t. $\rho_n(\mathcal{T})$. Then, there exists an \mathfrak{L}_n -interpretation \mathcal{I} that satisfies $\rho_n(\mathcal{T})$ and an $x \in \Delta^{\mathcal{I}}$ such that $(C^{n-1})^{\mathcal{I}}(x) > (D^{n-1})^{\mathcal{I}}(x)$. By Fact 1, $(C^{n-1})^{\mathcal{I}}(x) = 1$ and $(D^{n-1})^{\mathcal{I}}(x) = 0$. This in particular means that $x \in C^{\mathcal{I}_{cr}} \setminus D^{\mathcal{I}_{cr}}$, yielding the following.

Proposition 5. *If C is subsumed by D w.r.t. \mathcal{T} , then C^{n-1} is 1-subsumed by D^{n-1} w.r.t. $\rho_n(\mathcal{T})$.*

To prove the converse, we construct from a classical interpretation $\mathcal{I} = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}})$ the \mathfrak{L}_n -interpretation $\mathcal{I}_n = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}_n})$, where:

- $A^{\mathcal{I}_n}(x) := 1$ if $x \in A^{\mathcal{I}}$ and $A^{\mathcal{I}_n}(x) := \frac{n-2}{n-1}$ otherwise, for every $A \in \mathbf{N}_C$ and $x \in \Delta^{\mathcal{I}}$,
- $r^{\mathcal{I}_n}(x, y) := 1$ if $(x, y) \in r^{\mathcal{I}}$ and $r^{\mathcal{I}_n}(x, y) := \frac{n-2}{n-1}$ otherwise, for every $r \in \mathbf{N}_R$ and $x, y \in \Delta^{\mathcal{I}}$.

As before, this transformation preserves the satisfaction of the TBox w.r.t. the operator ρ_n .

Lemma 6. *If a classical interpretation \mathcal{I} satisfies \mathcal{T} , then \mathcal{I}_n satisfies $\rho_n(\mathcal{T})$.*

Proof Sketch. We again consider only two cases. Consider first any $\langle A \sqsubseteq (\exists r.B)^{n-1} \geq \frac{1}{n-1} \rangle \in \rho_n(\mathcal{T})$ and $x \in \Delta^{\mathcal{I}}$. If $((\exists r.B)^{n-1})^{\mathcal{I}_n}(x) = 0$, then

$$1 > (\exists r.B)^{\mathcal{I}_n}(x) = \sup_{z \in \Delta^{\mathcal{I}_n}} r^{\mathcal{I}_n}(x, z) *_{\mathfrak{L}_n} B^{\mathcal{I}_n}(z).$$

Therefore, every $y \in \Delta^{\mathcal{I}}$ must satisfy either $r^{\mathcal{I}_n}(x, y) < 1$ or $B^{\mathcal{I}_n}(y) < 1$. By the definition of \mathcal{I}_n , we get $x \notin (\exists r.B)^{\mathcal{I}}$, and thus $x \notin A^{\mathcal{I}}$ by assumption. Again by the definition of \mathcal{I}_n , we have $A^{\mathcal{I}_n}(x) = \frac{n-2}{n-1}$, and hence $A^{\mathcal{I}_n}(x) \Rightarrow_{\mathfrak{L}_n} ((\exists r.B)^{n-1})^{\mathcal{I}_n}(x) = \frac{1}{n-1}$.

In the case that $((\exists r.B)^{n-1})^{\mathcal{I}_n}(x) > 0$, Fact 1 yields that $((\exists r.B)^{n-1})^{\mathcal{I}_n}(x) = 1$ and

$$A^{\mathcal{I}_n}(x) \Rightarrow_{\mathfrak{L}_n} ((\exists r.B)^{n-1})^{\mathcal{I}_n}(x) = 1 \geq \frac{1}{n-1}.$$

Consider now any $\langle A \sqsubseteq B_1 \sqcap B_2 \geq \frac{n-2}{n-1} \rangle \in \rho_n(\mathcal{T})$. If $(B_1 \sqcap B_2)^{\mathcal{I}_n}(x) < \frac{n-2}{n-1}$, then by the definition of \mathcal{I}_n we have $B_1^{\mathcal{I}_n}(x) = B_2^{\mathcal{I}_n}(x) = \frac{n-2}{n-1}$, and thus $x \notin B_1^{\mathcal{I}} \cup B_2^{\mathcal{I}}$. Since \mathcal{I} satisfies \mathcal{T} , this implies $x \notin A^{\mathcal{I}}$. Again by the definition of \mathcal{I}_n , $A^{\mathcal{I}_n}(x) = \frac{n-2}{n-1}$. Since $(B_1 \sqcap B_2)^{\mathcal{I}_n}(x) = \frac{n-2}{n-1}$, we can conclude that $A^{\mathcal{I}_n}(x) \Rightarrow_{\mathfrak{L}_n} (B_1 \sqcap B_2)^{\mathcal{I}_n}(x) = \frac{n-2}{n-1}$.

In the case that $(B_1 \sqcap B_2)^{\mathcal{I}_n}(x) \geq \frac{n-2}{n-1}$, we also have

$$A^{\mathcal{I}_n}(x) \Rightarrow_{\mathfrak{L}_n} (B_1 \sqcap B_2)^{\mathcal{I}_n}(x) \geq (B_1 \sqcap B_2)^{\mathcal{I}_n}(x) \geq \frac{n-2}{n-1}. \quad \square$$

Using arguments analogous to those sketched above, we obtain the following proposition.

Proposition 7. *If C is not subsumed by D w.r.t. \mathcal{T} , then C^{n-1} is not 1-subsumed by D^{n-1} w.r.t. $\rho_n(\mathcal{T})$.*

We have thus reduced classical subsumption in \mathcal{ELU} to 1-subsumption in \mathfrak{L}_n - \mathcal{EL} . Since the former is EXPTIME-hard [Baader *et al.*, 2005], we obtain an EXPTIME lower bound for the complexity of the latter. An exponential-time algorithm for solving subsumption in the more expressive language \mathfrak{L}_n - \mathcal{ALC} , which provides a matching upper bound, was presented in [Borgwardt and Peñaloza, 2013].

Theorem 8. *For any $n \geq 3$, deciding ℓ -subsumption with respect to a TBox in \mathfrak{L}_n - \mathcal{EL} is EXPTIME-complete.*

4 Arbitrary Finite Chains

We now show that the complexity result from the previous section can be transferred to almost all logics of the form \mathbf{L} - \mathcal{EL} where \mathbf{L} is a finite chain. More precisely, subsumption in \mathbf{L} - \mathcal{EL} is EXPTIME-complete for all finite chains except those of the form \mathbf{G}_n . For the latter, this problem can be shown to be tractable.

As detailed in Section 2, any finite chain \mathbf{L} that is not of the form \mathbf{G}_n must contain a finite Łukasiewicz chain in an interval $[a, b]$ with at least three elements. We use this fact to reduce subsumption in \mathfrak{L}_n - \mathcal{EL} to subsumption in \mathbf{L} - \mathcal{EL} , where n is the cardinality of the interval $[a, b]$ in \mathbf{L} that is isomorphic to \mathfrak{L}_n . We extend the bijection $\lambda: [a, b] \rightarrow \mathfrak{L}_n$ to the chain \mathbf{L} :

- $\lambda(x) := 0$ if $x < a$ and
- $\lambda(x) := 1$ if $x > b$.

We also use the inverse $\lambda^{-1}: \mathfrak{L}_n \rightarrow 2^{\mathbf{L}}$ of this function, for which we have $\lambda^{-1}(0) = [0, a]$, and $\lambda^{-1}(1) = [b, 1]$. When it is clear from the context, we will also use λ^{-1} to denote the inverse of the original bijection; i.e. $\lambda^{-1}: \mathfrak{L}_n \rightarrow [a, b]$. As shown in the following lemma, these operators are compatible with all the operators that are relevant to fuzzy \mathcal{EL} .

Lemma 9. *1. For all $p, q \in \mathbf{L}$, we have*

- $\lambda(p *_{\mathbf{L}} q) = \lambda(p) *_{\mathfrak{L}_n} \lambda(q)$, and
- if $q \geq a$, then $\lambda(p \Rightarrow_{\mathbf{L}} q) = \lambda(p) \Rightarrow_{\mathfrak{L}_n} \lambda(q)$.

2. For all values $p, q \in \mathfrak{L}_n$, $p' \in \lambda^{-1}(p) \cap [a, 1]$, and $q' \in \lambda^{-1}(q) \cap [a, 1]$, we have

- $p' *_{\mathbf{L}} q' \in \lambda^{-1}(p *_{\mathfrak{L}_n} q) \cap [a, 1]$, and
- $p' \Rightarrow_{\mathbf{L}} q' \in \lambda^{-1}(p \Rightarrow_{\mathfrak{L}_n} q) \cap [a, 1]$.

We can now describe the reduction from \mathfrak{L}_n - \mathcal{EL} to \mathbf{L} - \mathcal{EL} . Let \mathcal{T} be an \mathfrak{L}_n - \mathcal{EL} TBox, $\ell \in \mathfrak{L}_n \setminus \{0\}$, and A, B two concept names for which we want to check whether A is ℓ -subsumed by B w.r.t. \mathcal{T} . We define the \mathbf{L} - \mathcal{EL} TBox \mathcal{T}' as follows.

$$\begin{aligned} \mathcal{T}' := & \{ \langle C \sqsubseteq D \geq \lambda^{-1}(p) \rangle \mid \langle C \sqsubseteq D \geq p \rangle \in \mathcal{T} \} \cup \\ & \{ \langle \top \sqsubseteq D \geq a \rangle \mid \langle C \sqsubseteq D \geq p \rangle \in \mathcal{T} \} \cup \\ & \{ \langle \top \sqsubseteq B \geq a \rangle \}. \end{aligned}$$

Restricting all right-hand side of GCIs in \mathcal{T} to have values $\geq a$ is necessary in light of Lemma 9.

We prove that if A is $\lambda^{-1}(\ell)$ -subsumed by B w.r.t. \mathcal{T}' , then A is ℓ -subsumed by B w.r.t. \mathcal{T} . Given an \mathfrak{L}_n -interpretation \mathcal{I} , we define the \mathbf{L} -interpretation $\mathcal{I}_L = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}_L})$ for all $C \in \mathbf{N}_C$, $r \in \mathbf{N}_R$, and $x, y \in \Delta^{\mathcal{I}}$ as follows:

- $C^{\mathcal{I}_L}(x) := \lambda^{-1}(C^{\mathcal{I}}(x))$ and
- $r^{\mathcal{I}_L}(x, y) := \lambda^{-1}(r^{\mathcal{I}}(x, y))$.

Using the properties from Lemma 9, it is easy to see that if \mathcal{I} satisfies \mathcal{T} , then \mathcal{I}_L satisfies \mathcal{T}' .

Suppose now that A is not ℓ -subsumed by B w.r.t. \mathcal{T} . Then there is an \mathfrak{L}_n -interpretation \mathcal{I} that satisfies \mathcal{T} , and an $x \in \Delta^{\mathcal{I}}$ such that $A^{\mathcal{I}}(x) \Rightarrow_{\mathfrak{L}_n} B^{\mathcal{I}}(x) < \ell$. But then also $A^{\mathcal{I}_L}(x) \Rightarrow_{\mathfrak{L}_n} B^{\mathcal{I}_L}(x) < \lambda^{-1}(\ell)$ since λ^{-1} is strictly monotone. Since \mathcal{I}_L is a model of \mathcal{T}' , this proves the claim.

Proposition 10. *If A is $\lambda^{-1}(\ell)$ -subsumed by B w.r.t. \mathcal{T}' , then A is ℓ -subsumed by B w.r.t. \mathcal{T} .*

Conversely, given an L-interpretation \mathcal{I} , we can construct the \mathfrak{L}_n -interpretation $\mathcal{I}_n = (\Delta^{\mathcal{I}}, \cdot^{\mathcal{I}_n})$ where

- $C^{\mathcal{I}_n}(x) := \lambda(C^{\mathcal{I}}(x))$ for all $C \in \mathbf{N}_C$ and $x \in \Delta^{\mathcal{I}}$, and
- $r^{\mathcal{I}_n}(x, y) := \lambda(r^{\mathcal{I}}(x, y))$ for all $r \in \mathbf{N}_R$ and $x, y \in \Delta^{\mathcal{I}}$.

Using arguments similar to those presented before, it is possible to prove that if \mathcal{I} satisfies \mathcal{T}' , then \mathcal{I}_n satisfies \mathcal{T} . Thus, subsumption w.r.t. \mathcal{T}' can be decided by checking the subsumption w.r.t. the original TBox \mathcal{T} .

Proposition 11. *If A is ℓ -subsumed by B w.r.t. \mathcal{T} , then A is $\lambda^{-1}(\ell)$ -subsumed by B w.r.t. \mathcal{T}' .*

This shows that subsumption in \mathfrak{L}_n - \mathcal{EL} is polynomially reducible to subsumption in L- \mathcal{EL} , for any finite chain L containing an interval \mathfrak{L}_n . By Theorem 8, the latter problem is EXPTIME-hard. A matching upper bound is also a consequence of the results from [Borgwardt and Peñaloza, 2013]. As discussed before, every finite chain L is either of the form \mathbf{G}_n or contains an interval isomorphic to \mathfrak{L}_n , for some n . Overall, this yields the desired complexity result.

Theorem 12. *Let L be a finite chain that is not of the form \mathbf{G}_n . Deciding ℓ -subsumption with respect to a TBox in L- \mathcal{EL} is EXPTIME-complete.*

Together with Proposition 3, we thus obtain a full characterization of the complexity of reasoning in fuzzy \mathcal{EL} over finite chains, depending on the t-norm that defines the semantics. If all elements of the chain are idempotent w.r.t. the t-norm, then subsumption can be decided in PTIME. Otherwise (i.e. if there is at least one non-idempotent element), this problem becomes EXPTIME-hard. In the following section, we show that the exponential lower bound holds also for infinite chains that contain a Łukasiewicz component.

5 The Infinite Łukasiewicz T-norm

We now consider the infinite chain $[0, 1]$, and show EXPTIME-hardness for deciding subsumption in L- \mathcal{EL} , for any t-norm that contains a Łukasiewicz component (Definition 2). As shown in [Borgwardt and Peñaloza, 2013], it suffices to prove this result for any t-norm that starts with the Łukasiewicz t-norm. Thus, for the rest of this section we consider a continuous t-norm that is isomorphic to the infinitely valued Łukasiewicz t-norm in the interval $[0, b]$ for some $b \in (0, 1]$.

To obtain the EXPTIME lower bound, we reduce subsumption in \mathcal{ELU} to subsumption in L- \mathcal{EL} . This reduction is very similar to the construction from Section 3. The main difference is that, in order to guarantee that the constructed L- \mathcal{EL} TBox can be used to decide the original \mathcal{ELU} subsumption problem, we need to restrict its models in such a way that all relevant concepts can only take the values $\frac{b}{2}$ or $\geq b$.

Given a concept C , let \mathcal{T}_C be the L- \mathcal{EL} TBox

$$\mathcal{T}_C := \{\langle C^2 \sqsubseteq C^3 \geq 1 \rangle, \langle \top \sqsubseteq C \geq \frac{b}{2} \rangle\}.$$

Every model \mathcal{I} of this TBox must satisfy $C^{\mathcal{I}}(x) \geq \frac{b}{2}$ for every $x \in \Delta^{\mathcal{I}}$ due to the second axiom. The first axiom additionally guarantees that $C^{\mathcal{I}}(x) \notin (\frac{b}{2}, b)$ holds: if $\frac{b}{2} < C^{\mathcal{I}}(x) < b$, then $(C^2)^{\mathcal{I}}(x) = C^{\mathcal{I}}(x) + C^{\mathcal{I}}(x) - b > 0$, and thus $(C^3)^{\mathcal{I}}(x) < (C^2)^{\mathcal{I}}(x)$, violating the axiom.

Similar to the reduction in Section 3, we will use the truth degree $\frac{b}{2} \in \mathbf{L}$ to stand for “false” in \mathcal{ELU} and any degree greater or equal to b to represent “true.” We define the function $\rho_{\mathfrak{L}}$ for every \mathcal{ELU} GCI in normal form (cf. Section 3):

$$\rho_{\mathfrak{L}}(A_1 \sqcap A_2 \sqsubseteq B) := \langle A_1 \sqcap A_2 \sqsubseteq B \geq b \rangle$$

$$\rho_{\mathfrak{L}}(\exists r. A \sqsubseteq B) := \langle \exists r. A \sqsubseteq B \geq b \rangle$$

$$\rho_{\mathfrak{L}}(A \sqsubseteq \exists r. B) := \langle A \sqsubseteq (\exists r. B)^2 \geq \frac{b}{2} \rangle$$

$$\rho_{\mathfrak{L}}(A \sqsubseteq B_1 \sqcup B_2) := \langle A \sqsubseteq B_1 \sqcap B_2 \geq \frac{b}{2} \rangle.$$

Given an \mathcal{ELU} TBox \mathcal{T} in normal form, let $\mathbf{AC}(\mathcal{T})$ be the set of all concept names and existential restrictions appearing in \mathcal{T} . We extend the mapping $\rho_{\mathfrak{L}}$ to \mathcal{ELU} TBoxes as follows:

$$\rho_{\mathfrak{L}}(\mathcal{T}) := \{\rho_{\mathfrak{L}}(C \sqsubseteq D) \mid C \sqsubseteq D \in \mathcal{T}\} \cup \bigcup_{C \in \mathbf{AC}(\mathcal{T})} \mathcal{T}_C.$$

Let now $A, B \in \mathbf{N}_C$. One can show that A is subsumed by B w.r.t. \mathcal{T} iff A is b -subsumed by B w.r.t. $\rho_{\mathfrak{L}}(\mathcal{T}) \cup \mathcal{T}_A \cup \mathcal{T}_B$. The proof follows the same ideas presented in Section 3. The TBoxes \mathcal{T}_C ensure that only three values are relevant for the models, and hence L behaves like \mathfrak{L}_3 on them.

From the previous arguments, we see that for any continuous chain L that starts with Łukasiewicz, subsumption in L- \mathcal{EL} is EXPTIME-hard. As shown in [Borgwardt and Peñaloza, 2013], if L is the ordinal sum of \mathbf{L}_1 and \mathbf{L}_2 over the intervals $[0, a]$ and $[a, 1]$, respectively, for some $a \in (0, 1)$, then subsumption in L- \mathcal{EL} is at least as hard as subsumption in \mathbf{L}_2 - \mathcal{EL} . Additionally, every chain L that contains a Łukasiewicz component can be described as such an ordinal sum, where \mathbf{L}_2 starts with Łukasiewicz. This means that the EXPTIME-hardness holds for all such continuous chains.

Theorem 13. *If L is defined using any continuous t-norm over $[0, 1]$ containing a Łukasiewicz component, then deciding ℓ -subsumption w.r.t. a TBox in L- \mathcal{EL} is EXPTIME-hard.*

This improves the CO-NP lower bound from [Borgwardt and Peñaloza, 2013]. It is unknown whether a similar lower bound holds for t-norms containing only product components. An upper bound is known only for G- \mathcal{EL} , where subsumption can be decided in PTIME [Mailis et al., 2012].

6 Conclusions

We have shown that reasoning in finitely valued extensions of fuzzy \mathcal{EL} becomes exponentially harder than in classical \mathcal{EL} even if only one additional truth value interpreted under Łukasiewicz semantics is considered. This provides the first example of a finitely valued DL that exhibits an increased complexity compared to the underlying classical DL. The same complexity lower bound holds for any infinitely valued t-norm over $[0, 1]$ that contains a Łukasiewicz component.

Although EXPTIME-complete, we believe that subsumption in finitely valued \mathcal{EL} can be solved more efficiently than by the algorithms developed for expressive finitely valued DLs [Borgwardt and Peñaloza, 2013; 2014]. We plan to look at adaptations of consequence-based algorithms for classical DLs [Baader *et al.*, 2005; Kazakov, 2009]. On the theoretical side, we will investigate whether other inexpressive DLs like \mathcal{FL}_0 [Baader, 1990] or *DL-Lite* [Calvanese *et al.*, 2005] also exhibit an increase in complexity under Łukasiewicz semantics. We also want to study the effect of the product semantics on the complexity of these logics.

Acknowledgments

This work was supported by DFG under grant BA 1122/17-1 (FuzzyDL) and the Cluster of Excellence ‘cfAED’; and by the ESF project “POST UP II” No. CZ.1.07/2.3.00/30.0041, co-financed by the European Social Fund and the state budget of the Czech Republic. The work was developed while R. Peñaloza was affiliated with TU Dresden and cfAED.

References

- [Baader and Peñaloza, 2011] Franz Baader and Rafael Peñaloza. On the undecidability of fuzzy description logics with GCIs and product t-norm. In *Proc. of the 8th Int. Symp. on Frontiers of Combining Systems (FroCoS’11)*, volume 6989 of *LNAI*, pages 55–70. Springer, 2011.
- [Baader *et al.*, 2005] Franz Baader, Sebastian Brandt, and Carsten Lutz. Pushing the \mathcal{EL} envelope. In *Proc. of the 19th Int. Joint Conf. on Artificial Intelligence (IJCAI’05)*, pages 364–369. Morgan-Kaufmann, 2005.
- [Baader, 1990] Franz Baader. Terminological cycles in KL-ONE-based knowledge representation languages. In *Proc. of the 8th Nat. Conf. on Artificial Intelligence (AAAI’90)*, pages 621–626. AAAI Press, 1990.
- [Bobillo and Straccia, 2011] Fernando Bobillo and Umberto Straccia. Reasoning with the finitely many-valued Łukasiewicz fuzzy description logic *SRQLQ*. *Inf. Sci.*, 181:758–778, 2011.
- [Borgwardt and Peñaloza, 2013] Stefan Borgwardt and Rafael Peñaloza. Positive subsumption in fuzzy \mathcal{EL} with general t-norms. In *Proc. of the 23rd Int. Joint Conf. on Artificial Intelligence (IJCAI’13)*, pages 789–795. AAAI Press, 2013.
- [Borgwardt and Peñaloza, 2013] Stefan Borgwardt and Rafael Peñaloza. The complexity of lattice-based fuzzy description logic. *J. Data Semant.*, 2(1):1–19, 2013.
- [Borgwardt and Peñaloza, 2014] Stefan Borgwardt and Rafael Peñaloza. Consistency reasoning in lattice-based fuzzy description logics. *Int. J. Approx. Reas.*, 55(9):1917–1938, 2014.
- [Borgwardt *et al.*, 2014] Stefan Borgwardt, Marco Cerami, and Rafael Peñaloza. Many-valued horn logic is hard. In *Proc. of the 1st Int. Workshop on Logics for Reasoning about Preferences, Uncertainty, and Vagueness (PRUV’14)*, volume 1205 of *CEUR*, pages 52–58, 2014.
- [Borgwardt *et al.*, 2015a] Stefan Borgwardt, Marco Cerami, and Rafael Peñaloza. Subsumption in finitely valued fuzzy \mathcal{EL} . LTCS-Report 15-06, TU Dresden, Germany, 2015. See <http://lat.inf.tu-dresden.de/research/reports.html>.
- [Borgwardt *et al.*, 2015b] Stefan Borgwardt, Felix Distel, and Rafael Peñaloza. The limits of decidability in fuzzy description logics with general concept inclusions. *Artif. Intell.*, 218:23–55, 2015.
- [Bou *et al.*, 2012] Felix Bou, Marco Cerami, and Francesc Esteva. Concept satisfiability in finite-valued fuzzy description logics is PSPACE-complete (extended abstract). In *Proc. of the 3rd Conf. on Logic, Algebras and Truth Degrees (LATD’12)*, pages 49–54, 2012.
- [Calvanese *et al.*, 2005] Diego Calvanese, Giuseppe De Giacomo, Domenico Lembo, Maurizio Lenzerini, and Riccardo Rosati. *DL-Lite*: Tractable description logics for ontologies. In *Proc. of the 20th Nat. Conf. on Artificial Intelligence (AAAI’05)*, pages 602–607. AAAI Press, 2005.
- [Cerami and Straccia, 2013] Marco Cerami and Umberto Straccia. On the (un)decidability of fuzzy description logics under Łukasiewicz t-norm. *Inf. Sci.*, 227:1–21, 2013.
- [Cerami and Straccia, 2014] Marco Cerami and Umberto Straccia. Complexity sources in fuzzy description logic. In *Proc. of the 27th Int. Workshop on Description Logics (DL’14)*, volume 1193 of *CEUR*, pages 421–433, 2014.
- [Hájek, 2001] Petr Hájek. *Metamathematics of Fuzzy Logic (Trends in Logic)*. Springer-Verlag, 2001.
- [Hájek, 2005] Petr Hájek. Making fuzzy description logic more general. *Fuzzy Sets Syst.*, 154(1):1–15, 2005.
- [Kazakov, 2009] Yevgeny Kazakov. Consequence-driven reasoning for Horn *SHIQ* ontologies. In *Proc. of the 21st Int. Joint Conf. on Artificial Intelligence (IJCAI’09)*, pages 2040–2045. AAAI Press, 2009.
- [Klement *et al.*, 2000] Erich Peter Klement, Radko Mesiar, and Endre Pap. *Triangular Norms*. Trends in Logic, Studia Logica Library. Springer-Verlag, 2000.
- [Mailis *et al.*, 2012] Theofilos Mailis, Giorgos Stoilos, Nikolaos Simou, Giorgos B. Stamou, and Stefanos Kollias. Tractable reasoning with vague knowledge using fuzzy \mathcal{EL}^{++} . *J. Intell. Inform. Syst.*, 39(2):399–440, 2012.
- [Mayor and Torrens, 1993] Gaspar Mayor and Joan Torrens. On a class of operators for expert systems. *Int. J. Intell. Syst.*, 8:771–778, 1993.
- [Mayor and Torrens, 2005] Gaspar Mayor and Joan Torrens. Triangular norms on discrete settings. In *Logical, Algebraic, Analytic and Probabilistic Aspects of Triangular Norms*, chapter 7, pages 189–230. Elsevier, 2005.
- [Mostert and Shields, 1957] Paul S. Mostert and Allen L. Shields. On the structure of semigroups on a compact manifold with boundary. *Ann. Math.*, 65(1):117–143, 1957.
- [Straccia, 2001] Umberto Straccia. Reasoning within fuzzy description logics. *Journal of Artificial Intelligence Research*, 14:137–166, 2001.