

Uncertainty, Belief, and Probability

Ronald Fagin and Joseph Y. Halpern
IBM Almaden Research Center, Dept. K53/S02
650 Harry Rd.
San Jose, CA 95120
email: fagin@ibm.com, halpern@ibm.com

Abstract

We introduce a new probabilistic approach to dealing with uncertainty, based on the observation that probability theory does not require that every event be assigned a probability. For a *nonmeasurable* event (one to which we do not assign a probability), we can talk about only the *inner measure* and *outer measure* of the event. Thus, the measure of belief in an event can be represented by an interval (defined by the inner and outer measure), rather than by a single number. Further, this approach allows us to assign a belief (inner measure) to an event E without committing to a belief about its negation $\neg E$ (since the inner measure of an event plus the inner measure of its negation is not necessarily one). Interestingly enough, inner measures induced by probability measures turn out to correspond in a precise sense to Dempster-Shafer belief functions. Hence, in addition to providing promising new conceptual tools for dealing with uncertainty, our approach shows that a key part of the important Dempster-Shafer theory of evidence is firmly rooted in classical probability theory.

1 Introduction

Dealing with uncertainty is a fundamental issue for AI. The most widely-used approach to dealing with uncertainty is undoubtedly the Bayesian approach. It has the advantage of relying on well-understood techniques from probability theory, as well as some philosophical justification on the grounds that a "rational" agent must assign uncertainties to events in a way that satisfies the axioms of probability [Cox46, Sav54]. On the other hand, the Bayesian approach has been widely criticized for requiring an agent to assign a subjective probability to every event. While this can be done in principle by having the agent play a suitable betting game [Jef83],¹ it does have a number of drawbacks. Among others, there is the computational difficulty of arriving at the probability. There is also the issue of whether it is reasonable

¹This idea is due to Ramsey [Ram31] and was rediscovered by von Neumann and Morgenstern [vNM47]; a clear exposition can be found in [LR57].

to describe confidence by a single point rather than a range. While an agent might be prepared to agree that the probability of an event lies within a given range, say between $1/3$ and $1/2$, he might not be prepared to say that it is precisely $.435$.

Not surprisingly, there has been a great deal of debate regarding the Bayesian approach (see [Che85] and [Sha76] for some of the arguments). Numerous other approaches to dealing with uncertainty have been proposed, including Dempster-Shafer theory [Dem68, Sha76], Cohen's model of endorsements [Coh85], and various nonstandard, modal, and fuzzy logics (for example, [HR87, Zad75]). A recent overview of the field can be found in [Saf88]. Of particular interest to us here is the Dempster-Shafer approach, which uses *belief functions* and *plausibility functions* to attach numerical lower and upper bounds on the likelihoods of events.

Although the Bayesian approach requires an agent to assign a probability to every event, probability theory does not. The usual reason that mathematicians deal with nonmeasurable events (those that are not assigned a probability) is out of mathematical necessity. For example, it is well known that if the sample space of the probability space consists of all numbers in the real interval $[0, 1]$, then we cannot allow every set to be measurable if (like Lebesgue measure) the measure is to be translation-invariant (see [Boy64, page 54]). However, in this paper we allow nonmeasurable events out of choice, rather than out of mathematical necessity. An event E for which an agent has insufficient information to assign a probability is modelled as a nonmeasurable set. The agent is not forced to assign a probability to E in our approach. We can provide meaningful lower and upper bounds on our degree of belief in E by using the standard mathematical notions of *inner measure* and *outer measure* induced by the probability measure [Hal50], which, roughly speaking, are the probability of the largest measurable event contained in E and the smallest measurable event containing E , respectively.

Allowing nonmeasurable events has its advantages. The uncertainty of event E is no longer given by a single number, but rather by an interval defined by the inner and outer measures. Furthermore, it is possible for the belief (i.e., inner measure) of event E to be a without the belief of $\neg E$ being $1 - a$. Rather than nonmeasurability being a mathematical nuisance, we have turned it

here into a desirable feature!

We feel that this paper makes three major contributions. The first is conceptual: In certain situations, our approach gives a useful way to think about and reason about uncertainty. In particular, the use of nonmeasurable sets seems to provide a useful way to capture our uncertainty about the probability of an event. The second is technical: We prove that, in a precise sense, inner measures induced by probability measures are equivalent to Shafer's belief functions (and so outer measures induced by probability measures are equivalent to Shafer's plausibility functions). The implications of this equivalence are significant. Although some, such as Cheeseman [Che85], consider the theory of belief functions as *ad hoc* and essentially nonprobabilistic (see discussion by Shafer [Sha86]), our results help show that a key part of the Dempster-Shafer theory of evidence is firmly rooted in classical probability theory. The last contribution is also technical: by combining our results here with those of a companion paper [FHM88], we are able to obtain a sound and complete axiomatization for a rich propositional logic of evidence, and provide a decision procedure for the satisfiability problem, which we show is no harder than that of propositional logic (NP-complete). Our techniques may provide a means for automatically deducing the consequences of a body of evidence.

2 Probability theory

To make our discussion precise, it is helpful to recall some basic definitions from probability theory (see [Fel57] for more details). A *probability space* (S, \mathcal{X}, μ) consists of a set S (called the *sample space*), a σ -algebra \mathcal{X} of subsets of S (i.e., a set of subsets of S containing S and closed under complementation and countable union, but not necessarily consisting of all subsets of S) whose elements are called *measurable sets*, and a *probability measure* $\mu: \mathcal{X} \rightarrow [0, 1]$ satisfying the following properties:

P1. $\mu(X) \geq 0$ for all $X \in \mathcal{X}$

P2. $\mu(S) = 1$

P3. $\mu(\bigcup_{i=1}^{\infty} X_i) = \sum_{i=1}^{\infty} \mu(X_i)$, if the X_i 's are pairwise disjoint members of \mathcal{X} .

Property P3 is called *countable additivity*. Of course, the fact that \mathcal{X} is closed under countable union guarantees that if each $X_i \in \mathcal{X}$, then so is $\bigcup_{i=1}^{\infty} X_i$. If \mathcal{X} is a finite set, then we can simplify property P3 above to

P3'. $\mu(X \cup Y) = \mu(X) + \mu(Y)$, if X and Y are disjoint members of \mathcal{X} .

This property is called *finite additivity*. Properties P1, P2, and P3' characterize probability measures in finite spaces.

In a probability space (S, \mathcal{X}, μ) , the probability measure μ is not defined on 2^S (the set of all subset of S), but only on \mathcal{X} . We can extend μ to 2^S in two standard ways, by defining functions μ_* and μ^* , traditionally called the *inner measure* and *outer measure induced by μ* [Hal50]. For an arbitrary subset $A \subseteq S$, we define

$$\begin{aligned} \mu_*(A) &= \sup\{\mu(X) \mid X \subseteq A \text{ and } X \in \mathcal{X}\} \\ \mu^*(A) &= \inf\{\mu(X) \mid X \supseteq A \text{ and } X \in \mathcal{X}\}. \end{aligned}$$

If there are only finitely many measurable sets (in particular, if S is finite), then it is easy to see that the inner measure of A is the measure of the largest measurable set contained in A , while the outer measure of A is the measure of the smallest measurable set containing A . In any case, it is not hard to show by countable additivity that for each set A , there are measurable sets B and C where $B \subseteq A \subseteq C$ such that $\mu(B) = \mu_*(A)$ and $\mu(C) = \mu^*(A)$. Note that if there are no nonempty measurable sets contained in A , then $\mu_*(A) = 0$, and if there are no measurable sets containing A other than the whole space S , then $\mu^*(A) = 1$. The properties of probability spaces guarantee that if X is a measurable set, then $\mu_*(X) = \mu^*(X) = \mu(X)$. In general we have $\mu^*(A) = 1 - \mu_*(\bar{A})$.

Suppose we have a situation we want to reason about. Typically we do so by fixing a finite set $\Phi = \{p_1, \dots, p_n\}$ of primitive propositions, which can be thought of as corresponding to basic events, such as "it is raining now" or "the coin landed heads". The set $\mathcal{L}(\Phi)$ of (*propositional*) *formulas* is the closure of Φ under the Boolean operations \wedge and \neg . The primitive propositions in Φ do not in general describe mutually exclusive events. To get mutually exclusive events, we can consider all the *atoms*, that is, all the formulas of the form $p'_1 \wedge \dots \wedge p'_n$, where p'_i is either p_i or $\neg p_i$. Let At denote the set of atoms.

We have been using the word "event" informally, sometimes meaning "set" and sometimes meaning "formula". We now want to be more formal, and to be able to talk explicitly about the probability of a *formula*. However, a probability measure is a function on sets, not formulas. Fortunately, it is easy to go from sets to formulas.

Using standard propositional reasoning, it is easy to see that any formula can be written as a disjunction of atoms. Thus, a formula φ can be identified with the unique set $\{\delta_1, \dots, \delta_k\}$ of atoms such that $\varphi \equiv \delta_1 \vee \dots \vee \delta_k$. If we want to assign probabilities to all formulas, we can simply assign probabilities to each of the atoms, and then use the finite additivity property of probability measures to compute the probability of an arbitrary formula. This amounts to taking a probability space of the form $(At, 2^{At}, \mu)$. The states in the probability space are just the atoms, and the measurable subsets are all the sets of atoms (i.e., all formulas). Once we assign a measure to the singleton sets (i.e., to the atoms), we can extend by additivity to any subset. We call such a probability space a *Nilsson structure*, since this is essentially what Nilsson used to give meaning to formulas in his probability logic [Nil86].² Given a Nilsson structure $N = (At, 2^{At}, \mu)$ and a formula φ , let $W_N(\varphi)$ denote the *weight* or *probability* of φ in N , which is defined to be $\mu(At(\varphi))$, where $At(\varphi)$ is the set of atoms whose disjunction is equivalent to φ .

A more general approach is to take a *probability structure* to be a tuple $(S, \mathcal{X}, \mu, \pi)$, where (S, \mathcal{X}, μ) is a probability space, and π associates with each $s \in S$ a truth assignment $\pi(s): \Phi \rightarrow \{\mathbf{true}, \mathbf{false}\}$. We say that p is

² Actually, the use of possible worlds in giving semantics to probability formulas goes back to Carnap [Car50].

true at s if $\pi(s)(p) = \text{true}$; otherwise, we say that p is false at s .

We think of S as consisting of the possible states of the world. We can associate with each state s in S a unique atom describing the truth values of the primitive propositions in s . For example, if $\Phi = \{p_1, p_2\}$, and if $\pi(s)(p_1) = \text{true}$ and $\pi(s)(p_2) = \text{false}$, then we associate with s the atom $p_1 \wedge \neg p_2$. It is perfectly all right for there to be several states associated with the same atom (indeed, there may be an infinite number, since we allow S to be infinite, even though Φ is finite). This situation may occur if a state is not completely characterized by the events that are true there. This is the case, for example, if there are features of worlds that are not captured by the primitive propositions.

We can easily extend $\pi(s)$ to a truth assignment on all formulas by taking the usual rules of propositional logic. Then if M is a probability structure, we can associate with every formula φ the set φ^M consisting of all the states in M where φ is true (i.e., the set $\{s \in S \mid \pi(s)(\varphi) = \text{true}\}$). Of course, we assume that n is defined so that $\text{true}^M = S$. If p^M is measurable for every primitive proposition $p \in \Phi$, then φ^M is also measurable for every formula φ (since the set X of measurable sets is closed under complementation and countable union). We say M is a *measurable probability structure* if φ^M is measurable for every formula φ .

It makes sense to talk about the probability of $\langle p \rangle$ in M only if φ^M is measurable; we can then take the probability of φ , which we denote $W_M(\varphi)$, to be $\mu(\varphi^M)$. If φ^M is not measurable, then we cannot talk about its probability. However, we can still talk about its inner measure and outer measure, since these are defined for all subsets. Intuitively, the inner and outer measure provide lower and upper bounds on the probability of φ . In general, if φ^M is not measurable, then we take $W_M(\varphi)$ to be $\mu_*(\varphi)$, i.e., the inner measure of φ in M .

We define a probability structure M and a Nilsson structure N to be *equivalent* if $W_M(\varphi) = W_N(\varphi)$ for every formula φ . Intuitively, a probability structure and a Nilsson structure are equivalent if they assign the same probability to every formula. The next theorem shows that there is a natural correspondence between Nilsson structures and measurable probability structures.³

Theorem 2.1:

1. For every Nilsson structure there is an equivalent measurable probability structure.
2. For every measurable probability structure there is an equivalent Nilsson structure.

Why should we even allow nonmeasurable sets? As the following example shows (as do others given in the full paper), using nonmeasurability allows us to avoid assigning probabilities to those events for which we have insufficient information to assign a probability.

Example 2.2: Ron has two blue suits and two gray suits. He has a very simple method for deciding what color suit to wear on any particular day: he simply

The proof of this and all other theorems mentioned here can be found in the full paper [FH88],

tosses a (fair) coin: if it lands heads he wears a blue suit, and if it lands tails he wears a gray suit. Once he's decided what color suit to wear, he just chooses the rightmost suit of that color on the rack. Both of Ron's blue suits are single-breasted, while one of Ron's gray suits is single-breasted and the other is double-breasted. Ron's wife Susan is (fortunately for Ron) a little more fashion-conscious than he is. She also knows how Ron makes his sartorial choices. So, from time to time, she makes sure that the gray suit she considers preferable is to the right (which it depends on current fashions and perhaps on other whims of Susan).⁴ Suppose we don't know about the current fashions (or about Susan's current whims). What can we say about the probability of Ron's wearing a single-breasted suit on Monday?

In terms of possible worlds, it is clear that there are four possible worlds, one corresponding to each of the suits that Ron could choose. For definiteness, suppose states s_1 and s_2 correspond to the two blue suits, s_3 corresponds to the single-breasted gray suit, and s_4 corresponds to the double-breasted gray suit. Let $S = \{s_1, s_2, s_3, s_4\}$. There are two features of interest about a suit: its color and whether it is single-breasted or double-breasted. Let the primitive proposition g denote "the suit is gray" and let db denote "the suit is double-breasted", and define the truth assignment n in the obvious way. Note that the atom $\neg g \wedge \neg db$ is associated with both states s_1 and s_2 . Since the two blue suits are both single-breasted, these two states cannot be distinguished by the formulas in our language.

What are the measurable events? Besides S itself and the empty set, the only other candidates are $\{s_1, s_2\}$ ("Ron chooses a blue suit") and $\{s_3, s_4\}$ ("Ron chooses a gray suit"). However, $SB = \{s_1, s_2, s_3\}$ ("Ron chooses a single-breasted suit") is nonmeasurable. The reason is that we do not have a probability on the event "Ron chooses a single-breasted suit, given that Ron chooses a gray suit", since this in turn depends on the probability that Susan put the single-breasted suit to the right of the other gray suit, which we do not know. Susan's choice might be characterizable by a probability distribution; it might also be deterministic, based on some complex algorithm which even she might not be able to describe; or it might be completely nondeterministic, in which case it is not technically meaningful to talk about the "probability" of Susan's actions! Our ignorance here is captured by nonmeasurability. Informally, we can say that the probability of Ron choosing a single-breasted suit lies somewhere in the interval $[1/2, 1]$, since it is bounded below by the probability of Ron choosing a blue suit. This is an informal statement because formally it does not make sense to talk about the probability of a nonmeasurable event. The formal analogue is simply that the inner measure of SB is $1/2$, while its outer measure is 1 .

⁴Any similarity between the characters in this example and the first author of this paper and his wife Susan is not totally accidental.

3 The Dempster-Shafer theory of evidence

The Dempster-Shafer theory of evidence [Sha76] provides another approach to attaching likelihoods to events. This theory starts out with a *belief function* (sometimes called a *support function*). For every event (i.e., set) A , the belief in A , denoted $Bel(A)$, is a number in the interval $[0,1]$ that places a lower bound on likelihood of A . We have a corresponding number $Pl(A) = 1 - Bel(A)$, called the *plausibility* of A , which places an upper bound on the likelihood of A . Thus, to every event A we can attach the interval $[Bel(A), Pl(A)]$. Like a probability measure, a belief function assigns a "weight" to subsets of a set S , but unlike a probability measure, the domain of a belief function is always taken to be *all* subsets of S . Just as we defined probability structures, we can define a *DS structure* (where, of course, "DS" stands for Dempster-Shafer) to be a tuple (S, Bel, TT) , where S and TT are as before, and where $Bel: 2^S \rightarrow [0, 1]$ is a function satisfying:

B1. $Bel(\emptyset) = 0$

B2. $Bel(S) = 1$

B3. $Bel(A_1 \cup \dots \cup A_k) \geq \sum_{I \subseteq \{1, \dots, k\}, I \neq \emptyset} (-1)^{|I|+1} Bel(\bigcap_{i \in I} A_i)$.

A belief function is typically defined on a *frame of discernment*, consisting of mutually exclusive and exhaustive propositions describing the domain of interest. We think of the set S of states in a belief structure as being this frame of discernment. We could always choose S to be some subset of At , the set of atoms, so that its elements are in fact propositions in the language. In general, given a DS structure $D = (S, Bel, \pi)$ and formula φ , we define the weight $W_D(\varphi)$ to be $Bel(\varphi^D)$, where φ^D is the set of states where φ is true. Thus we can talk about an agent's degree of belief in $\langle \varphi \rangle$ in D , described by $W_D(\varphi)$, by identifying $\langle \varphi \rangle$ with the set φ^D and considering the belief in φ^D . As before, we define a probability structure M (resp., a Nilsson structure N , a DS structure D') and a DS structure D to be *equivalent* if $WM(\varphi) = W_D(\varphi)$ (resp., $WN(\varphi) = W_D(\varphi)$, $W_{D'}(\varphi) = W_D(\varphi)$) for every formula φ .

Property B3 may seem unmotivated. Perhaps the best way to understand it is as an analogue to the usual inclusion-exclusion rule for probabilities [Fel57, p. 89], which is obtained by replacing the inequality by equality (and the belief function Bel by a probability measure). In particular, B3 is important for probability measures (we prove a more general result, namely that it holds for all inner measures induced by probability measures, in Proposition 3.1 below). Hence, if (S, X, μ) is a probability space and $X = 2^S$ (making every subset of S measurable), then μ is a belief function. (This fact has been observed frequently before; see, for example, [Sha76].) It follows that every Nilsson structure is a DS structure.

It is easy to see that the converse does not hold. For example, suppose there is only one primitive proposition, say p , in the language, so that $At = \{p, \neg p\}$, and let $D_0 = (At, Bel, \pi)$ be such that $Bel(\{p\}) = 1/2$, $Bel(\{\neg p\}) = 0$, and π is defined in the obvious way.

Intuitively, there is weight of evidence $1/2$ for p , and no evidence for $\neg p$. Thus $W_{D_0}(p) = 1/2$ and $W_{D_0}(\neg p) = 0$. D_0 is not equivalent to any Nilsson structure, since if N is a Nilsson structure such that $WN(p) = 1/2$, then we must have $WN(\neg p) = 1/2$.

These observations tell us that in some sense belief functions are more general than probability measures, provided we restrict attention to probability spaces where all sets are measurable. This fact is well known. Indeed, in [Sha76], Shafer makes explicit use of the greater generality of belief functions. While he does consider events E such that $Bel(\neg E) = 1 - Bel(E)$ (he calls such events *probabilistic*), he also wants to allow non-probabilistic events. He gives examples of events where the fact that we would like to assign weight $.8$ to our belief in event E does not mean that we want to assign weight $.2$ to our belief in $\neg E$. In our framework, where we allow nonmeasurable sets, we can view probabilistic events as corresponding to measurable sets, while non-probabilistic events do not. We can push this analogy much further. Not only do nonmeasurable sets correspond to non-probabilistic events, but the inner measures induced by probability measures correspond to belief functions.

Proposition 3.1: // (S, X, p) is a probability space, then f_{μ} is a belief function on 2^S .

Proposition 3.1 says that every inner measure is a belief function (and thus generalizes the statement that every probability measure is a belief function). The converse does not quite hold. For example, consider the DS space D_0 defined above. There is no probability measure μ that we can define on $\{p, \neg p\}$ such that $\mu_* = Bel$. However, it is easy to define a probability structure M such that $\mu_*(p^M) = 1/2$ and $\mu_*(\neg p^M) = 0$. That is, we can find a probability structure equivalent to D_0 . The next theorem generalizes this observation.

Theorem 3.2:

1. For every DS structure there is an equivalent probability structure.
2. For every probability structure there is an equivalent DS structure.

Intuitively, Theorem 3.2 says that belief functions and inner measures induced by probability measures are precisely the same if their domains are considered to be formulas rather than sets. As we shall see, this result has important implications regarding complete axiomatizations and decision procedures.

4 Reasoning about belief and probability

We are often interested in the inferences we can make about probabilities or beliefs given some information. In order to do this, we need a language for doing such reasoning. Such a language is given in [FHM88]. A *term* in this language is an expression of the form $a_1 w(\varphi_1) + \dots + a_k w(\varphi_k)$, where a_1, \dots, a_k are integers and $\varphi_1, \dots, \varphi_k$ are propositional formulas. A *basic weight formula* is one of the form $t \geq b$, where t is a

term and b is an integer. A *weight formula* is a Boolean combination of basic weight formulas. We sometimes use obvious abbreviations without further comment, such as $w(\varphi) \geq w(\psi)$ for $w(\varphi) - w(\psi) \geq 0$.

We give semantics to the formulas in our language with respect to all the structures we have been considering. Let A' be either a Nilsson structure, a probability structure, or a DS structure, and let l be a weight formula. We now define what it means for K to *satisfy* l , written $K \models l$. For a basic weight formula,

$$K \models a_1 w(\varphi_1) + \dots + a_k w(\varphi_k) \geq b \text{ iff} \\ a_1 W_K(\varphi_1) + \dots + a_k W_K(\varphi_k) \geq b.$$

We then extend \models in the obvious way to conjunctions and negations. The interpretation of $w(\varphi)$ is either "the probability of φ " (for Nilsson structures or measurable probability structures), "the inner measure of φ " for general probability structures, or "the belief in φ " (for DS structures).

Let K be a class of structures (in the cases of interest to us, K is the class of either probability structures, measurable probability structures, Nilsson structures, or DS structures). As usual, we define a weight formula l to be *satisfiable with respect to* K if $K \models l$ for some $K \in \mathcal{K}$. Similarly, l is *valid with respect to* K if $K \models l$ for all $K \in \mathcal{K}$.

In [FHM88], an axiom system *AX ME AS FOR* reasoning about measurable probability structures is provided. The system has three parts, which deal respectively with propositional reasoning, reasoning about linear inequalities, and reasoning about probability. For example, a typical axiom for reasoning about linear inequalities is

$$(a_1 w(\varphi_1) + \dots + a_k w(\varphi_k) \geq b) \Rightarrow \\ (ca_1 w(\varphi_1) + \dots + ca_k w(\varphi_k) \geq cb) \text{ if } c \geq 0,$$

which says that both sides of an inequality can be multiplied by a positive constant. (The remaining axioms for reasoning about inequalities are described in the full paper.)

For reasoning about probability, we have the following axioms. The first three correspond to the usual laws of probability, except that W3 corresponds to finite additivity, not countable additivity.

W1. $w(\varphi) \geq 0$ (nonnegativity)

W2. $w(\text{true}) = 1$ (the probability of the event *true* is 1)

W3. $w(\varphi \wedge \psi) + w(\varphi \wedge \neg\psi) = w(\varphi)$ (additivity)

W4. $w(\varphi) = w(\psi)$ if $\varphi \equiv \psi$ is a propositional tautology

As is shown in [FHM88], *AXMEAS* characterizes the valid formulas for measurable probability structures.

Theorem 4.1: ([FHM88]) *AX_{MEAS} is a sound and complete axiomatization for weight formulas with respect to measurable probability structures.*

This result, together with Theorem 2.1, immediately gives us

Corollary 4.2: *AXMEAS is a sound and complete axiomatization for weight formulas with respect to Nilsson structures.*

AX MEAS is not sound with respect to arbitrary probability structures, where $w(\varphi)$ is interpreted as the inner measure of φ^M . In particular, axiom W3 no longer holds: inner measures are not finitely additive. Let *AX* be obtained from *AXMEAS* by replacing W3 by the following two axioms, which are obtained from conditions B1 and B3 for belief functions in an obvious way:

W5. $w(\text{false}) = 0$

W6. $w(\varphi_1 \vee \dots \vee \varphi_k) \geq \sum_{I \subseteq \{1, \dots, k\}, I \neq \emptyset} (-1)^{|I|+1} w(\bigwedge_{i \in I} \varphi_i)$

Theorem 4.3: ([FHM88]) *AX is a sound and complete axiomatization for weight formulas with respect to probability structures.*

Applying Theorem 3.2, we immediately get

Corollary 4.4: *AX is a sound and complete axiomatization for weight formulas with respect to DS structures.*

Thus, using *AX*, we can derive all consequences of a collection of beliefs.

Combining the preceding results with results of [FHM88], we can also characterize the complexity of reasoning about probability and belief.

Theorem 4.5: *The complexity of deciding whether a weight formula is satisfiable with respect to probability structures (respectively, measurable probability structures, Nilsson structures, DS structures) is NP-complete.*

(This result in the case of Nilsson structures was obtained independently in [GKP88].) Note that Theorem 4.5 says that reasoning about probability and belief is, in a precise sense, exactly as difficult as propositional reasoning. This is the best we could expect, since it is easy to see that reasoning about probability and belief is at least as hard as propositional reasoning (the propositional formula φ is satisfiable iff the weight formula $w(\varphi) > 0$ is satisfiable).

5 Combining evidence

An important issue for belief functions, each of which can be viewed as representing a distinct body of evidence, is how to combine them to obtain a new belief function that somehow reflects the combined evidence. A way of doing so is provided by Dempster's *rule of combination*, which was introduced by Dempster [Dem68] and was further developed and studied in an elegant and rather complete manner by Shafer [Sha76].

In the full paper [FH88], we show that there is a natural way (in the spirit of Dempster's rule) to define the combination $D_x \oplus D_2$ of two DS structures D_1 and D_2 , and a natural way to define the combination $M_1 \otimes M_2$ of two probability structures M_1 and M_2 , such that the following theorem holds.

Theorem 5.1: *Let D_1 and D_2 be DS structures. There are probability structures M_1 and M_2 such that (a) D_1 is equivalent to M_1 , (b) D_2 is equivalent to M_2 , and (c) $D_1 \oplus D_2$ is equivalent to $M_1 \otimes M_2$.*

This theorem shows that the spirit of Dempster's rule of combination can be captured within probability theory. We are currently investigating alternatives

to Dempster's rule of combination for revising beliefs about uncertainty in the presence of new information. The idea is to consider what it means to take a conditional probability with respect to a nonmeasurable set. We plan to report on this work in a future paper.

6 Related work

Although we believe we are the first to propose using inner and outer measures as a way of dealing with uncertainty, there are a number of other works with similar themes. We briefly discuss them here.

A number of authors have argued that we should think in terms of an interval in which the probability lies, rather than a unique numerical probability (see, for example, [Kyb61, Kyb88]). Good [Goo62], Koopman [Koo40a, Koo40b], and Smith [Smi61] try to derive reasonable properties for the intuitive notions of *lower* and *upper* probability, which are somehow meant to capture lower and upper bounds on an agent's belief in a proposition. Good observes that "The analogy [between lower and upper probability and] inner and outer measure is obvious. But the axioms for upper and lower probability do not all follow⁷ from the theory of inner and outer measure."

Dempster [Dem66, Dem68] gives a formal mathematical definition of lower and upper probability in terms of a tuple (S, μ, T, Γ) , which we call a *Dempster structure*. $(S, 2^S, \mu)$ is a probability space (Dempster assumes for simplicity that every subset of S is measurable). T is another set, and Γ is a multi-valued mapping S to T . Thus, $\Gamma(s)$ is a subset of T for each $s \in S$. Given $A \subseteq T$, we define subsets A_* and A^* of S as follows:

$$\begin{aligned} A_* &= \{s \in S \mid \Gamma(s) \neq \emptyset, \Gamma(s) \subseteq A\} \\ A^* &= \{s \in S \mid \Gamma(s) \cap A \neq \emptyset\}. \end{aligned}$$

Provided $\mu(T^*) \neq 0$, we define the lower and upper probabilities of A , written $P_*(A)$ and $P^*(A)$ respectively, by

$$\begin{aligned} P_*(A) &= \mu(A_*)/\mu(T^*) \\ P^*(A) &= \mu(A^*)/\mu(T^*). \end{aligned}$$

It is easy to check that $T_* = T^*$. Thus, dividing by $\mu(T^*)$ has the effect of normalizing so that $P_*(T) = P^*(T) = 1$.

There is a close relationship between lower and upper probabilities and inner and outer measures induced by a probability measure. Given a probability structure $M = (S, \mathcal{X}, \mu, \pi)$ where S is finite, let $(\mathcal{X}', \mu', T, \Gamma)$ be the Dempster structure where (1) \mathcal{X}' is a basis for \mathcal{X} ,⁵ (2) μ' is a probability measure defined on $2^{\mathcal{X}'}$ by taking $\mu'(\{A\}) = \mu(A)$ for $A \in \mathcal{X}'$ and then extending to all subsets of \mathcal{X}' by finite additivity, (3) T consists of

⁵A subset X' of X is said to be a *basis* (of X) if the members of X' are nonempty and disjoint, and if X consists precisely of countable unions of members of X' . It is easy to see that if X is finite then it has a basis. Moreover, whenever X has a basis, it is unique: it consists precisely of the minimal elements of X (the nonempty sets none of whose nonempty subsets are in X). Note that if X has a basis, once we know the probability of every set in the basis, we can compute the probability of every measurable set by using countable additivity.

all propositional formulas, and (4) for $A \in \mathcal{X}'$, we define $F(A)$ to consist of all formulas φ such that φ is true at some point in A (in the structure M). Thus T is a multivalued mapping from \mathcal{X}' to T . It is easy to check that $P_*(\{\varphi\}) = \mu_*(\varphi^M)$ and $P^*(\{\varphi\}) = \mu^*(\varphi^M)$ for all formulas φ .

Ruspini [Rus87] also considers giving semantics to probability formulas by using possible worlds, but he includes epistemic notions in the picture. Briefly, his approach can be described as follows (where we have taken the liberty of converting some of his notation to ours, to make the ideas easier to compare). Fix a set $\{p_1, \dots, p_n\}$ of primitive propositions. Instead of considering just propositional formulas, Ruspini allows epistemic formulas; he obtains his language by closing off under the propositional connectives $\wedge, \vee, \Rightarrow$, and \neg as well as the epistemic operator K . Thus, a typical formula in his language would be $K(p_1 \Rightarrow K(p_2 \wedge p_3))$. (A formula such as $K\varphi$ should be read "the agent knows φ ".) Rather than considering arbitrary sample spaces as we have done here, where at each point in the sample space some subset of primitive propositions is true, Ruspini considers one fixed sample space S (which he calls a *sentence space*) whose points consist of all the possible truth assignments to these formulas consistent with the axioms of the modal logic S5. (See, for example, [HM85] for an introduction to S5. We remark that it can be shown that there are less than 2^{2^2} consistent truth assignments, so that S is finite.) We can define an equivalence relation \sim on S by taking $s \sim t$ if s and t agree on the truth values of all formulas of the form $K\varphi$. The equivalence classes form a basis for a σ -algebra of measurable subsets of S . Let \mathcal{X} be this σ -algebra. For any formula φ , let φ^S consist of all the truth assignments in S that make φ true. It is easy to check that $(K\varphi)^S$, the set of truth assignments that make $K\varphi$ true, is the union of equivalence classes, and hence is measurable. Let μ be any probability measure defined on \mathcal{X} . Given s , we can consider the probability structure $(S, \mathcal{X}, \mu, \pi)$, where we take $\pi(s)(p) = s(p)$. (Since s is a truth assignment, this is well defined.) The axioms of S5 guarantee us that $\{F \mid \langle p \rangle^S\}$ is the largest measurable subset contained in φ^M ; thus, $\mu_*(\varphi^M) = \mu((K\varphi)^S)$.

Ruspini then considers the DS structure (At, Bel, π') , where π' is defined in the obvious way on the atoms in At , and $Bel(\varphi^D) = \mu((K\varphi)^S) (= \mu_*(\varphi^M))$.⁶ Ruspini shows that Bel defined in this way is indeed a belief function. Thus, Ruspini shows a close connection between probabilities, inner measures, and belief functions in the particular structures that he considers. He does not show a general relationship between inner measures and belief functions; in particular, he does not show that DS structures are equivalent to probability structures, as we do in Theorem 3.2.

In the full paper, we explore further relations between our work and that of Ruspini, as well as comparing our characterization of belief functions with those of Shafer [Sha79], Kyburg [Kyb87], and Pearl [Pea88].

⁶Ruspini actually defines the belief function directly on formulas; i.e., he defines $Bel(\varphi)$. In our notation, what he is doing is defining a weight function WD -

Acknowledgments

The authors are grateful to Glenn Shafer, Moshe Vardi, and Henry Kyburg for helpful comments and suggestions.

References

- [Car50] R. Carnap, *Logical Foundations of Probability*, University of Chicago Press, 1950.
- [Che85] P. Cheeseman, In defense of probability, *Ninth International Joint Conference on Artificial Intelligence (IJCAI-85)*, 1985, pp. 1002-1009.
- [Coh85] P. R. Cohen, *Heuristic Reasoning about Uncertainty: An Artificial Intelligence Approach*, Pitman, 1985.
- [Cox46] R. Cox, Probability, frequency, and reasonable expectation, *American Journal of Physics* 14:1, 1946, pp. 1-13.
- [Dem66] A. P. Dempster, Upper and lower probabilities induced by a multivalued mapping, *Annals of Mathematical Statistics* 38, 19(36), pp. 325-339.
- [Dem68] A. P. Dempster, A generalization of Bayesian inference, *Journal of the Royal Statistical Society, Series B* 30, 1968, pp. 205-247.
- [Fel57] W. Feller, *An Introduction to Probability Theory and its Applications*, Vol. I, John Wiley & Sons, 2nd edition, 1957.
- [FH88] R. Fagin and J. Y. Halpern, Uncertainty, belief, and probability, IBM Research Report RJ 6191, 1988.
- [FIIM88] R. Fagin, J. Y. Halpern, and N. Megiddo, A logic for reasoning about probabilities, *Proc. 3rd IEEE Symp. on Logic in Computer Science*, 1988, pp. 277-291. A revised and expanded version of this paper appears as IBM Research Report RJ 6190, April 1988.
- [GKP88] G. Georgakopoulos, D. Kavvadias, and C. H. Papadimitriou, Probabilistic satisfiability, *Journal of Complexity* 4:1, 1988, pp. 1-11.
- [Goo62] I. J. Good, The measure of a non-measurable set, *Logic, Methodology, and Philosophy of Science* (E. Nagel, P. Suppes, and A. Tarski, eds.), Stanford University Press, 1962, pp. 319-329.
- [Hal50] P. Halmos, *Measure Theory*, Van Nostrand, 1950.
- [HM85] J. Y. Halpern and Y. Moses, A guide to the modal logics of knowledge and belief, *Ninth International Joint Conference on Artificial Intelligence (IJCAI-85)*, 1985, pp. 480-490.
- [HR87] J. Y. Halpern and M. O. Rabin, A logic to reason about likelihood, *Artificial Intelligence* 32:3, 1987, pp. 379-405.
- [Jef83] R. C. Jeffrey, *The Logic of Decision*, University of Chicago Press, 1983.
- [Koo40a] B. O. Koopman, The axioms and algebra of intuitive probability, *Annals of Mathematics* 41, 1940, pp. 269-292.
- [Koo40b] B. O. Koopman, The bases of probability, *Bulletin of the American Mathematical Society* 46, 1940, pp. 763-774.
- [Kyb61] H. E. Kyburg, Jr., *Probability and the Logic of Rational Belief* Wesleyan University Press, 1961.
- Kyb87] H. E. Kyburg, Jr., Bayesian and non-Bayesian evidential updating, *Artificial Intelligence* 31, 1987, pp. 271-293.
- [Kyb88] H. E. Kyburg, Jr., Higher order probabilities and intervals, *International Journal of Approximate Reasoning* 2, 1988, pp. 195-209.
- [LR.57] R. D. Luce and H. Raiffa, *Games and Decisions*, Wiley, 1957.
- [Nil86] N. Nilsson, Probabilistic logic, *Artificial Intelligence* 28, 1986, pp. 71-87.
- [Pea88] J. Pearl, *Probabilistic Reasoning in Intelligent Systems*, Morgan Kaufmann, 1988.
- [Ram31] F. P. Ramsey, Truth and probability, *The Foundations of Probability and other Logical Essays* (R. B. Braithwaite, ed.), Harcourt Brace, 1931.
- [Roy64] H. L. Royden, *Real Analysis*, Macmillan, 1964.
- [Rus87] E. H. Ruspini, *The logical foundations of evidential reasoning*, Technical Report Technical Note 408, revised version, SRI International, 1987.
- Saf88] A. SafTiotti, An AI view of the treatment of uncertainty, *The Knowledge Engineering Review* 2:2, 1988.
- Sav54] L. J. Savage, *Foundations of Statistics*, John Wiley & Sons, 1954.
- Sha76] G. Shafer, *A Mathematical Theory of Evidence*, Princeton University Press, 1976.
- Sha79] G. Shafer, Allocations of probability, *Annals of Probability* 7:5, 1979, pp. 827-839.
- Sha86] (J. Shafer, The combination of evidence, *International Journal of Intelligent Systems* 1, 1986, pp. 155-179.
- Smi61] C. A. B. Smith, Consistency in statistical inference and decision, *Journal of the Royal Statistical Society, Series B* 23, 1961, pp. 1-25.
- [vNM47] J. von Neumann and O. Morgenstern, *Theory of Games and Economic Behavior*, Princeton University Press, 2nd edition, 1947.
- Zad75] L. A. Zadeh, Fuzzy logics and approximate reasoning, *Synthese* 30, 1975, pp. 407-428.